

Nature Medicine 2018 vol.24 N11, pages 1677-1682

Neuromodulation of lumbosacral spinal networks enables independent stepping after complete paraplegia

Gill M., Grahn P., Calvert J., Linde M., Lavrov I., Strommen J., Beck L., Sayenko D., Van Straaten M., Drubach D., Veith D., Thoreson A., Lopez C., Gerasimenko Y., Edgerton V., Lee K., Zhao K.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018, The Author(s), under exclusive licence to Springer Nature America, Inc. Spinal sensorimotor networks that are functionally disconnected from the brain because of spinal cord injury (SCI) can be facilitated via epidural electrical stimulation (EES) to restore robust, coordinated motor activity in humans with paralysis^{1–3}. Previously, we reported a clinical case of complete sensorimotor paralysis of the lower extremities in which EES restored the ability to stand and the ability to control step-like activity while side-lying or suspended vertically in a body-weight support system (BWS)⁴. Since then, dynamic task-specific training in the presence of EES, termed multimodal rehabilitation (MMR), was performed for 43 weeks and resulted in bilateral stepping on a treadmill, independent from trainer assistance or BWS. Additionally, MMR enabled independent stepping over ground while using a front-wheeled walker with trainer assistance at the hips to maintain balance. Furthermore, MMR engaged sensorimotor networks to achieve dynamic performance of standing and stepping. To our knowledge, this is the first report of independent stepping enabled by task-specific training in the presence of EES by a human with complete loss of lower extremity sensorimotor function due to SCI.

<http://dx.doi.org/10.1038/s41591-018-0175-7>

References

- [1] Harkema, S. et al. Effect of epidural stimulation of the lumbosacral spinal cord on voluntary movement, standing, and assisted stepping after motor complete paraplegia: a case study. *Lancet* 377, 1938–1947 (2011).
- [2] Angeli, C. A., Edgerton, V. R., Gerasimenko, Y. P. & Harkema, S. J. Altering spinal cord excitability enables voluntary movements after chronic complete paralysis in humans. *Brain* 137, 1394–1409 (2014).
- [3] van den Brand, R. et al. Restoring voluntary control of locomotion after paralyzing spinal cord injury. *Science* 336, 1182–1185 (2012).
- [4] Grahn, P. J. et al. Enabling task-specific volitional motor functions via spinal cord neuromodulation in a human with paraplegia. *Mayo Clin. Proc.* 92, 544–554 (2017).
- [5] Gerasimenko, Y. P. et al. Epidural spinal cord stimulation plus quipazine administration enable stepping in complete spinal adult rats. *J. Neurophysiol.* 98, 2525–2536 (2007).
- [6] Ichiyama, R. M., Gerasimenko, Y. P., Zhong, H., Roy, R. R. & Edgerton, V. R. Hindlimb stepping movements in complete spinal rats induced by epidural spinal cord stimulation. *Neurosci. Lett.* 383, 339–344 (2005).
- [7] Lavrov, I. et al. Epidural stimulation induced modulation of spinal locomotor networks in adult spinal rats. *J. Neurosci.* 28, 6022–6029 (2008).
- [8] Capogrosso, M. et al. A brain-spine interface alleviating gait deficits after spinal cord injury in primates. *Nature* 539, 284–288 (2016).

- [9] Wenger, N. et al. Spatiotemporal neuromodulation therapies engaging muscle synergies improve motor control after spinal cord injury. *Nat. Med.* 22, 138–145 (2016).
- [10] Wenger, N. et al. Closed-loop neuromodulation of spinal sensorimotor circuits controls refined locomotion after complete spinal cord injury. *Sci. Transl. Med.* 6, 255ra133 (2014).
- [11] Lavrov, I. et al. Facilitation of stepping with epidural stimulation in spinal rats: role of sensory input. *J. Neurosci.* 28, 7774–7780 (2008).
- [12] Dimitrijevic, M. R., Gerasimenko, Y. & Pinter, M. M. Evidence for a spinal central pattern generator in humans. *Ann. NY Acad. Sci.* 860, 360–376 (1998).
- [13] Minassian, K. et al. Stepping-like movements in humans with complete spinal cord injury induced by epidural stimulation of the lumbar cord: electromyographic study of compound muscle action potentials. *Spinal Cord* 42, 401–416 (2004).
- [14] Danner, S. M. et al. Human spinal locomotor control is based on flexibly organized burst generators. *Brain* 138, 577–588 (2015).
- [15] Rattay, F., Minassian, K. & Dimitrijevic, M. R. Epidural electrical stimulation of posterior structures of the human lumbosacral cord: 2. quantitative analysis by computer modeling. *Spinal Cord* 38, 473–489 (2000).
- [16] Kakulas, B. A. Pathology of spinal injuries. *Cent. Nerv. Syst. Trauma* 1, 117–129 (1984).
- [17] Dimitrijevic, M. R., Dimitrijevic, M. M., McKay, W. B. & Sherwood, A. M. EMG evidence of suprasegmental influence on motor unit activity in paralyzed muscles. *Clin. Neurophysiol.* 56, S68 (1983).
- [18] Dimitrijević, M. R. Residual motor functions in spinal cord injury. *Adv. Neurol.* 47, 138–155 (1988).
- [19] Taccola, G., Sayenko, D., Gad, P., Gerasimenko, Y. & Edgerton, V. R. And yet it moves: recovery of volitional control after spinal cord injury. *Prog. Neurobiol.* 160, 64–81 (2018).
- [20] Dimitrijevic, M. R. et al. Human spinal cord motor control that is partially or completely disconnected from the brain. *Am. J. Neuroprot. Neuroregen.* 8, 12–26 (2016).
- [21] Minassian, K. & Hofstoetter, U. S. Spinal cord stimulation and augmentative control strategies for leg movement after spinal paralysis in humans. *CNS Neurosci. Ther.* 22, 262–270 (2016).
- [22] Sherwood, A. M., Dimitrijevic, M. R. & McKay, W. B. Evidence of subclinical brain influence in clinically complete spinal cord injury: discomplete SCI. *J. Neurol. Sci.* 110, 90–98 (1992).
- [23] Field-Fote, E. C. & Roach, K. E. Influence of a locomotor training approach on walking speed and distance in people with chronic spinal cord injury: a randomized clinical trial. *Phys. Ther.* 91, 48–60 (2011).
- [24] Harkema, S. J., Schmidt-Read, M., Lorenz, D. J., Edgerton, V. R. & Behrman, A. L. Balance and ambulation improvements in individuals with chronic incomplete spinal cord injury using locomotor training-based rehabilitation. *Arch. Phys. Med. Rehabil.* 93, 1508–1517 (2012).
- [25] Forrest, G. F. et al. Neuromotor and musculoskeletal responses to locomotor training for an individual with chronic motor complete AIS-B spinal cord injury. *J. Spinal Cord Med.* 31, 509–521 (2008).
- [26] Dobkin, B. et al. Weight-supported treadmill vs over-ground training for walking after acute incomplete SCI. *Neurology* 66, 484–493 (2006).
- [27] Rejc, E., Angeli, C. A., Bryant, N. & Harkema, S. J. Effects of stand and step training with epidural stimulation on motor function for standing in chronic complete paraplegics. *J. Neurotrauma* 34, 1787–1802 (2017).
- [28] Carhart, M. R., He, J., Herman, R., D'Luzansky, S. & Willis, W. T. Epidural spinal-cord stimulation facilitates recovery of functional walking following incomplete spinal-cord injury. *IEEE Trans. Neural. Syst. Rehabil. Eng.* 12, 32–42 (2004).
- [29] Rejc, E., Angeli, C. & Harkema, S. Effects of lumbosacral spinal cord epidural stimulation for standing after chronic complete paralysis in humans. *PLoS One* 10, e0133998 (2015).
- [30] Gerasimenko, Y. P. et al. Noninvasive reactivation of motor descending control after paralysis. *J. Neurotrauma* 32, 1968–1980 (2015).
- [31] Lu, D. C. et al. Engaging cervical spinal cord networks to reenact volitional control of hand function in tetraplegic patients. *Neurorehabil. Neural Repair.* 30, 951–962 (2016).
- [32] Gad, P. et al. Weight bearing over-ground stepping in an exoskeleton with non-invasive spinal cord neuromodulation after motor complete paraplegia. *Front. Neurosci.* 11, 333 (2017).
- [33] Huang, H., He, J., Herman, R. & Carhart, M. R. Modulation effects of epidural spinal cord stimulation on muscle activities during walking. *IEEE Trans. Neural. Syst. Rehabil. Eng.* 14, 14–23 (2006).
- [34] Shah, P. K. & Lavrov, I. Spinal epidural stimulation strategies: clinical implications of locomotor studies in spinal rats. *Neuroscientist* 23, 664–680 (2017).
- [35] Courtine, G. & Bloch, J. Defining ecological strategies in neuroprosthetics. *Neuron* 86, 29–33 (2015).
- [36] Moritz, C. T. Now is the critical time for engineered neuroplasticity. *Neurotherapeutics* 15, 628–634 (2018).
- [37] Shah, P. K. et al. Variability in step training enhances locomotor recovery after a spinal cord injury. *Eur. J. Neurosci.* 36, 2054–2062 (2012).

- [38] Behrman, A. L. & Harkema, S. J. Locomotor training after human spinal cord injury: a series of case studies. *Phys. Ther.* 80, 688-700 (2000).
- [39] Harkema, S. J., Behrman, A. L. & Barbeau, H. in *Locomotor Training: Principles and Practice*, 54-84 (Oxford University Press, Oxford, UK, 2011).
- [40] Sayenko, D. G., Angeli, C., Harkema, S. J., Edgerton, V. R. & Gerasimenko, Y. P. Neuromodulation of evoked muscle potentials induced by epidural spinal-cord stimulation in paralyzed individuals. *J. Neurophysiol.* 111, 1088-1099 (2014).
- [41] Sayenko, D. G. et al. Spinal segment-specific transcutaneous stimulation differentially shapes activation pattern among motor pools in humans. *J. Appl. Physiol.* 118, 1364-1374 (2015).
- [42] Sayenko, D. G. et al. Effects of paired transcutaneous electrical stimulation delivered at single and dual sites over lumbosacral spinal cord. *Neurosci. Lett.* 609, 229-234 (2015).